

Seizures Nausea Headache Muscle c
d reflexes Lethargy Vomiting Gait diso
s Gait disorder Depressed reflexes Vor
Seizures Confusion Disorientation An
Muscle cramps Dizziness Anorexia Letha
ait disorder Nausea Vomiting Gait diso
Muscle cramps Restlessness Nausea D
sed reflexes Seizures Disorientation D
norexia Muscle cramps Confusion Re
Lethargy Drowsiness Restlessness N
sea Headache Vomiting Depressed re
ressed reflexes Gait disorder Lethargy
Nausea Disorientation Seizures Anore

Read between the lines.

Hyponatraemia secondary to SIADH.

The symptoms of hyponatraemia are often attributed to other conditions.¹
Without prompt treatment symptoms can worsen and put patients' health at risk.²

(SIADH is syndrome of inappropriate antidiuretic hormone secretion)

To find out more visit www.hyponatraemia.com



Otsuka-people creating new products for better health worldwide

References: 1. Miller M. *J Am Geriatr Soc* 2006; 54(2): 345–353.
2. Adrogué H & Madias NE. *N Engl J Med* 2000; 342(21): 1581–1589.

Date of preparation: January 2012 OPEL/0112/SAM/1013



Read between the lines. Hyponatraemia secondary to SIADH.

The symptoms of hyponatraemia are often attributed to other conditions.¹
Without prompt treatment symptoms can worsen and put patients' health at risk.²

(SIADH is syndrome of inappropriate antidiuretic hormone secretion)

To find out more visit www.hyponatraemia.com



Read between the lines.
Hyponatraemia secondary to SIADH.

The symptoms of hyponatraemia are often attributed to other conditions.¹
Without prompt treatment symptoms can worsen and put patients' health at risk.²

(SIADH is syndrome of inappropriate antidiuretic hormone secretion)

To find out more visit www.hyponatraemia.com

